

## Appendix

---

### $p$ -adic Numbers

In Chap. 7 we saw that the moduli space of curves over the complex numbers comes in a natural manner with an arithmetic structure. Just to remind you let me repeat the following facts. If  $\mathcal{C}$  is a curve, then it is the set of common zeros of finitely many homogeneous polynomials. For example, a curve in the projective plane can be given by the zeros of one homogeneous polynomial. If after a suitable change of coordinates the polynomials can be given in such a way that all coefficients are rational numbers then we say that the curve  $\mathcal{C}$  can be *defined over the rational numbers*  $\mathbb{Q}$ .

If  $g$  is a rational polynomial of degree  $k$  we can write it as

$$g = \sum_i r_i m_i, \quad r_i = \frac{s_i}{t_i}$$

where  $m_i$  are monomials (just products of the variables) and the  $s_i$  and  $t_i$  are integers. If we now multiply  $g$  by the least common multiple  $t$  of the  $t_i$  we get

$$\hat{g} = t \cdot g = \sum_i \hat{s}_i \cdot m_i$$

with  $\hat{s}_i$  integers. Of course  $g$  and  $\hat{g}$  have the same set of points as zeros. Hence every curve defined over the rational numbers  $\mathbb{Q}$  can be defined over the integers  $\mathbb{Z}$ . Hence we can use all number theoretic techniques to get information on the geometric (and arithmetic) structure. (This works also in the opposite direction.)

Now, how do we get from the rational numbers to the complex numbers? Of course this is well known. But the technique is also essential in the case of the  $p$ -adic numbers, so let me repeat it briefly. For the rational numbers we have the so-called *absolute value*. It allows us to define a topology and the notion of a convergent sequence. As you know there exist sequences of rational numbers which “converge” but have no limits in  $\mathbb{Q}$  (of course strictly speaking we do not talk about convergence in this case). We enlarge  $\mathbb{Q}$  by adding these limit points to get the real numbers  $\mathbb{R}$ .

One way to construct  $\mathbb{R}$  in more precise terms is the following. We call a sequence  $(x_n)$  a Cauchy sequence if for every rational number  $\epsilon > 0$  there exists a natural number  $n_0$  such that

$$|x_n - x_m| < \epsilon \quad \text{if } n, m > n_0.$$

As remarked above not every Cauchy sequence has to have a limit. But if it has a limit and this limit is 0, then we call  $(x_n)$  a *zero sequence*.

We can define addition, subtraction and multiplication on the set of Cauchy sequences element-wise, for example

$$(x_n) + (y_n) := (x_n + y_n).$$

The real numbers are now defined as the set of classes of Cauchy sequences modulo zero sequences. This means that we identify two Cauchy sequences  $(x_n)$  and  $(z_n)$  if  $(x_n) - (z_n)$  is a zero sequence. All operations can be well defined on these classes. In addition if  $[(x_n)]$  is not the zero class (equivalently  $(x_n)$  is not a zero sequence) we can define the multiplicative inverse  $[(x_n)]^{-1}$  in the following way. By adding a suitable zero sequence to  $(x_n)$  we can arrange that for the new sequence (still denoted by  $(x_n)$ )  $x_n \neq 0$  for all  $n$ . In this way we stay in the same class. We set

$$[(x_n)]^{-1} := \left[ \left( \frac{1}{x_n} \right) \right]$$

which is again a Cauchy sequence, because the  $x_n$  are bounded away from 0.

It is easy to see that  $\mathbb{R}$  will be a field and that  $\mathbb{Q}$  is embedded as a subfield via the constant sequence

$$[(x_n)] \quad \text{for all } n : x_n = x \in \mathbb{Q}.$$

Now we can also define the absolute value in  $\mathbb{R}$  by setting

$$|[(x_n)]| := [|x_n|].$$

With this we see that in  $\mathbb{R}$  all Cauchy sequences have a limit. Hence we call  $\mathbb{R}$  the completion of  $\mathbb{Q}$  with respect to the absolute value  $|\cdot|$ .

From the mathematical viewpoint  $\mathbb{R}$  is not yet fully satisfying: we cannot solve every algebraic equation. In technical terms  $\mathbb{R}$  is not *algebraically closed*. For this reason we have to “add” a single element, the root  $i$  of the polynomial equation

$$x^2 + 1 = 0.$$

It obeys the multiplicative law  $i^2 = -1$ . In this manner we get the complex numbers  $\mathbb{C}$ . We can extend the absolute value in  $\mathbb{R}$  to the usual complex absolute value.  $\mathbb{C}$  is now an algebraically closed complete field.

But what forces us to stick to this special absolute value to start from the rational numbers? Let us list the characteristic features of the absolute value which allowed us to work as above. The absolute value is a map

$$|\cdot| : \mathbb{Q} \rightarrow \mathbb{Q}$$

with

$$|a| \geq 0, \quad a \in \mathbb{Q}, \quad |a| = 0 \quad \text{if and only if} \quad a = 0. \quad (1)$$

$$|a \cdot b| = |a| \cdot |b|, \quad a, b \in \mathbb{Q} \quad (\text{multiplicative law}) \quad (2)$$

$$|a + b| \leq |a| + |b|, \quad a, b \in \mathbb{Q} \quad (\text{triangle identity}) \quad (3)$$

If we have such a map we call it a *valuation* for the field  $\mathbb{Q}$ . We can generalize this immediately to arbitrary fields (for the values we remain in  $\mathbb{Q}$  or  $\mathbb{R}$ ).

There is one map, the so-called *trivial valuation*, defined by

$$|a| = \begin{cases} 1, & \text{if } a \neq 0 \\ 0, & \text{if } a = 0. \end{cases}$$

This valuation is of no interest to us. If we use the word valuation we always assume a nontrivial one.

Let us consider a very interesting one. If

$$r = \frac{n}{m} \in \mathbb{Q}, \quad r \neq 0, \quad n, m \in \mathbb{Z}$$

then we can write

$$n = 2^s n', \quad m = 2^t m'$$

where  $n'$  and  $m'$  are integers which are not divisible by 2. We set

$$|r|_2 := 2^{t-s} \quad \text{and} \quad |0|_2 := 0.$$

Let us check the conditions above.

Condition 1: is clear.

Condition 2: Normalized as above we get

$$\begin{aligned} r_1 &= \frac{n_1}{m_1} = 2^l \cdot \frac{n'_1}{m'_1}, & r_2 &= \frac{n_2}{m_2} = 2^p \cdot \frac{n'_2}{m'_2}, \\ |r_1|_2 \cdot |r_2|_2 &= 2^{-l} 2^{-p} = 2^{-(l+p)} \\ r_1 r_2 &= 2^{l+p} \cdot \frac{n'_1 n'_2}{m'_1 m'_2} \end{aligned}$$

Because 2 is a prime number it divides neither  $n'_1 n'_2$  nor  $m'_1 m'_2$ . Hence we get

$$|r_1 r_2|_2 = 2^{-(l+p)}.$$

This is what we had to show.

Condition 3: We keep the same notation as above. We separate two cases. ( $l \neq p$ ): without restriction we assume  $l < p$ .

$$r_1 + r_2 = 2^l \cdot \left( \frac{n'_1}{m'_1} + 2^{p-l} \frac{n'_2}{m'_2} \right) = 2^l \cdot \frac{n'_1 m'_2 + 2^{p-l} n'_2 m'_1}{m'_1 m'_2}.$$

Now 2 divides neither the numerator (because 2 divides only one term in the sum) nor the denominator, hence

$$|r_1 + r_2|_2 = 2^{-l} = |r_1|_2 \leq |r_1|_2 + |r_2|_2.$$

( $l = p$ ): here we get

$$r_1 + r_2 = 2^l \cdot \frac{n'_1 m'_2 + n'_2 m'_1}{m'_1 m'_2}.$$

It can happen that 2 divides the numerator, hence

$$|r_1 + r_2|_2 = 2^{-(l+q)} \leq |r_1|_2 \leq |r_1|_2 + |r_2|_2$$

with  $q$  a positive integer.

In both cases condition 3 is fulfilled. Even more is true. We can replace (3) by

$$|r_1 + r_2| \leq \max(|r_1|, |r_2|) \quad (3')$$

and we know in addition

$$|r_1 + r_2| = \max(|r_1|, |r_2|) \quad \text{if } |r_1| \neq |r_2|$$

If the stronger condition (3') is valid we call  $|\cdot|$  a *nonarchimedian* valuation. Another name for it is an *ultrametric*. In a nonarchimedian valuation the integers are always within the “unit circle”. We can see this as follows. We have  $|1| = 1$  as usual ( $|a| = |1 \cdot a| = |1| \cdot |a|$ ). But now

$$|2| = |1 + 1| \leq \max(|1|, |1|) = |1| = 1.$$

By induction we get  $|n| \leq 1$  for all integers  $n$ . This is a rather strange result in contrast to the usual absolute value.

In the above the only fact we used about the number 2 was that it was a prime number. Hence we can define in an identical manner

$$|a|_p = p^{-s} \quad \text{if } a = p^s \cdot \frac{n'}{m'}, \quad |0|_p = 0$$

where  $p$  divides neither  $n'$  nor  $m'$ . We call  $|\cdot|_p$  the *p-adic valuation*. If there is a danger of confusion we use  $|\cdot|_\infty$  for the usual absolute value.

We can define a topology, convergence, Cauchy sequences and so on. With this we see that all these valuations are essentially different. To see this let us consider the sequence

$$(x_n) = (p^n)$$

for a fixed prime number  $p$ . Now

$$|p^n|_\infty = p^n$$

and hence it is unbounded and diverges in the usual topology. On the other hand

$$|p^n|_p = p^{-n}, \quad |p^n|_q = 1, \quad \text{for } q \neq p.$$

We see  $(x_n)$  is a zero sequence in the  $p$ -adic topology and a bounded (but not zero) sequence in the  $q$ -adic topology.


We call two valuations  $|\cdot|_a, |\cdot|_b$  equivalent if we have

$$|\cdot|_a = |\cdot|_b^r$$

with a positive real number  $r$ , i.e. if they define the same topology.

**Theorem A.1.** (*Ostrowski*)  $|\cdot|_\infty$  and  $|\cdot|_p$ , ( $p$  all prime numbers), are representatives of all equivalence classes of valuations for the rational number field  $\mathbb{Q}$ .

In the same way as we constructed the real numbers  $\mathbb{R}$  via Cauchy sequences with respect to the absolute value we construct the  $p$ -adic numbers  $\mathbb{Q}_p$  starting with the  $p$ -adic valuation for every prime number  $p$ . Our field  $\mathbb{Q}$  lies in all these completions:



In  $\mathbb{Q}_p$  we can do analysis similar to real analysis. Just as we can represent every real number as a decimal expansion, we can represent every  $p$ -adic number by a  $p$ -adic expansion. Take  $a \in \mathbb{Q}_p$ , by extension of the  $p$ -adic valuation to  $\mathbb{Q}_p$  we calculate

$$|a|_p = p^{-m}, \quad m \in \mathbb{Z}.$$

With this we can write

$$a = \sum_{k \geq m} a_k p^k = \lim_{n \rightarrow \infty} \sum_{k \geq m}^n a_k p^k$$

with

$$a_k \in \{0, 1, 2, \dots, p-1\}.$$

As you can see the sum goes in the opposite direction of the decimal expansion.

In some sense the analysis in  $\mathbb{Q}^p$  is much easier than that in  $\mathbb{R}$  but more unfamiliar. For example we have in  $\mathbb{Q}^p$  the following:

**Lemma A.2.** *A series  $\sum_k a_k$  converges if and only if the sequence  $(a_k)$  is a zero sequence.*

(Quite a number of students starting to learn mathematics would be happy if this would work also in  $\mathbb{R}$ .)

*Proof.* Let  $(a_k)$  be a zero sequence. Then  $(m \geq n)$

$$\left| \sum_{k=1}^m a_k - \sum_{k=1}^n a_k \right| = \left| \sum_{k=n}^m a_k \right| \leq \max_{k=n}^m |a_k|.$$

Because  $(a_k)$  is a zero sequence the right-hand side can be made as small as one wants by choosing  $n$  big enough. This means that the sequence of partial sums is a Cauchy sequence and hence has a limit. The other direction is the usual argument of real analysis.  $\square$

Now we can also define  $p$ -adic power series like the *exponential*

$$E(x) := \sum_{k=0}^{\infty} \frac{x^k}{k!}$$

and the *logarithm*

$$\log(1+x) = \sum_{k=1}^{\infty} (-1)^{k+1} \frac{x^k}{k}.$$

Despite the fact that the expressions are the same you might expect that the domain of convergence will be different. We write

$$|x|_p = p^{-\text{ord}_p(x)}$$

to define the  $p$ -adic order. With this we can calculate that  $E(x)$  is convergent if and only if  $\text{ord}_p(x) > \frac{1}{p-1}$  and  $\log(1+x)$  is convergent if and only if  $\text{ord}_p(x) > 0$ .

This is again a rather strange behaviour. If  $p \neq 2$  and  $x$  is an integer then  $E(x)$  is a convergent series if and only if  $x$  is a multiple of  $p$ . If  $p = 2$  and everything else stays the same, then  $x$  has to be a multiple of 4.

One can now introduce  $p$ -adic measures, distributions, integrals and Fourier transformations.<sup>1</sup> Let me mention that in  $\mathbb{Q}_p$  there is in contrast to  $\mathbb{R}$  still a lot of number theory involved. For example, we have also the  $p$ -adic integers which are defined to be the  $x \in \mathbb{Q}_p$  with  $\text{ord}_p(x) \geq 0$ . Like in the case of the rational numbers we can write every element of  $\mathbb{Q}_p$  as quotient of two  $p$ -adic integers.

We are still not at the end in our construction of the analogue of  $\mathbb{C}$ . To get from  $\mathbb{R}$  to  $\mathbb{C}$  we took the algebraic closure of  $\mathbb{R}$ . This can also be done with  $\mathbb{Q}_p$ . In this case we have to “add” infinitely many roots of polynomials to get

<sup>1</sup> See Koblitz, [Ko], p. 30.

to  $\mathbb{Q}_p^{a.c.}$ . Of course this  $\mathbb{Q}_p^{a.c.}$  also admits an extension of the  $p$ -adic valuation. But in contrast to  $\mathbb{C}$  it is no longer complete. We have to again construct the completion of this field. The resulting field is algebraically closed and complete.

Of course, one might ask why one should bother about the  $p$ -adic numbers if one is not interested in number theory. Some idea of their importance outside of number theory might be given by the following product formula for all rational numbers  $x \neq 0$ :

$$|x|_\infty \cdot \prod_{p \in \mathbb{P}} |x|_p = 1.$$

(We use  $\mathbb{P}$  to denote the set of all prime numbers.) It says, if we know all  $p$ -adic values of a rational number we also know its absolute value. This is a reformulation of the trivial fact that we know the absolute value of a rational number if we know how often all primes appear in the numerator and in the denominator. But there is more behind it. The mathematical idea is that the rational numbers are the objects of primary interest. But they are very difficult to handle from the arithmetic viewpoint.  $\mathbb{R}$  and all  $\mathbb{Q}_p$  are much easier. Each of them reflects one facet of the complexity of  $\mathbb{Q}$ . From this viewpoint one calls  $\mathbb{Q}$  a *global* field and  $\mathbb{R}$  and the different  $\mathbb{Q}_p$  *local* fields. If  $x \in \mathbb{Q}$  is a number fulfilling a suitable relation (for example to be a zero of a polynomial equation with integral coefficients) then  $x$  considered as element in  $\mathbb{R}$  and  $\mathbb{Q}_p$  clearly fulfils the transformed relation in  $\mathbb{R}$ , resp. in  $\mathbb{Q}_p$ . The converse problem is the one that really matters. If we can solve the transformed relations by elements  $x_\infty$  and  $x_p$ ,  $p \in \mathbb{P}$  (which in most cases is easier) we can ask under which conditions we can assemble this information to get a solution for the starting relation in  $\mathbb{Q}$ . (This is also known under the name *local-global principle*.)

In the above you might get the (right) feeling that it is better to consider the real number field and all  $p$ -adic number fields simultaneously. This is formalized in the concept of *adèles*. We start with the infinite product

$$\mathbb{R} \times \prod_{p \in \mathbb{P}} \mathbb{Q}_p.$$

It consists of all infinite sequences

$$(x_\infty, x_2, x_3, x_5, \dots, x_p, \dots)$$

where  $x_\infty \in \mathbb{R}, x_p \in \mathbb{Q}_p$ . Now we consider the subset with

$$|x|_p \leq 1$$

for almost every  $p$  (which is shorthand for: the condition can be violated only by a finite number of primes). This subset is called the set of *adèles* and is denoted by  $\mathbb{A}_\mathbb{Q}$ . It is a ring if we define addition and multiplication component-wise. The set of multiplicative invertible elements in  $\mathbb{A}_\mathbb{Q}$  is called *idèles*.  $\mathbb{Q}$  is now embedded diagonally into  $\mathbb{A}_\mathbb{Q}$  by

$$x \mapsto (x, x, x, \dots).$$

At first sight this is only a formal tool. But because this is a ring we can do algebraic geometry with it and other nice things. The following is an example taken from Manin (details can be found there [Man]). One can define the group  $\mathrm{Sl}(2, \mathbb{A}_{\mathbb{Q}})$  in a suitable manner. By diagonal embedding,  $\mathrm{Sl}(2, \mathbb{Q})$  is a discrete subgroup. Now the factor group is a compact group on which we can integrate. By normalizing the measure and splitting it up into the factors corresponding to factor groups of  $\mathrm{Sl}(2, \mathbb{R})$  and  $\mathrm{Sl}(2, \mathbb{Q}_p)$  we get the result

$$1 = \frac{\pi^2}{6} \cdot \prod_{p \in \mathbb{P}} (1 - p^{-2}).$$

This relates the arithmetically defined product expression with the transcendental number  $\pi$ . The product is an object which is defined by knowing the measure on the  $p$ -adic groups and it determines the analogous measure on the real group. Of course the above formula is not a new result gained only by the use of adèles. But the adèles gives a new insight. Classically it comes from the *Riemann zeta function*

$$\zeta(s) = \sum_{n \geq 1} \frac{1}{n^s}$$

where we have

$$\zeta(2) = \frac{\pi^2}{6} = \prod_{p \in \mathbb{P}} (1 - p^{-2})^{-1}$$

which is the above relation.

Up to now we have done everything over the rational numbers. In fact this is not sufficient. We have to allow finite algebraic field extensions of  $\mathbb{Q}$ . We can obtain these in the following way. We embed  $\mathbb{Q}$  in  $\mathbb{C}$  and take an  $\alpha \in \mathbb{C}$  which is a root of a rational irreducible polynomial  $f$ . (Here irreducible means it does not have a factorization into two nonconstant rational polynomials.) The imaginary unit  $i$  for example is such an  $\alpha$ . It is the zero of the polynomial

$$\mathbf{X}^2 + 1.$$

Another example is

$$\alpha = \zeta_3 = \exp\left(\frac{2\pi i}{3}\right),$$

a so-called 3rd root of unity. It is a zero of the polynomial

$$\mathbf{X}^2 + \mathbf{X} + 1.$$

We denote the smallest subfield of  $\mathbb{C}$  containing  $\mathbb{Q}$  and  $\alpha$  by  $\mathbb{Q}(\alpha)$ . It is also a finite-dimensional  $\mathbb{Q}$ -vector space. Its basis is given by

$$1, \alpha, \alpha^2, \dots, \alpha^m, \quad m = \deg f - 1.$$



Such a field is called a number field. Just as we can build  $\mathbb{Q}$  out of the integers as their quotients, we can find “integers” in  $\mathbb{Q}(\alpha)$  which do the job here. In contrast to the usual integers, the *rational integers*, these are called *algebraic integers*. With these algebraic integers we can do arithmetic, define divisibility and so on. Unfortunately we have in general no unique factorization of every algebraic integer into primes. Even worse, we have to drop the notion of prime number at all. But there is a rather useful substitute for the prime numbers. The prime numbers of  $\mathbb{Z}$  are in a 1:1 correspondence with the classes of nonarchimedean valuations of  $\mathbb{Q}$ . Now we ask for such nonarchimedean valuations for  $\mathbb{Q}(\alpha)$ . Let  $|\cdot|$  be such a valuation. By restricting it to  $\mathbb{Q}$  we get one of the  $p$ -adic valuations. Conversely one can show that for every  $p$ -adic valuation on  $\mathbb{Q}$ , there exist only finitely many extensions to the whole field  $\mathbb{Q}(\alpha)$ . By embedding into  $\mathbb{C}$  the absolute value is already extended to the usual complex absolute value on  $\mathbb{Q}(\alpha)$ . Different extensions of the absolute value correspond to different embeddings. There are also finitely many of them.

Now we can do everything (completion, adèles, ...) we did for  $\mathbb{Q}$  for this number field  $\mathbb{Q}(\alpha)$ . The corresponding complete field is a more tractable subfield of the big  $p$ -adic algebraically closed and complete field.

## Hints for Further Reading

Further details can be found in

[Ko] Koblitz, N.,  *$p$ -adic Numbers,  $p$ -adic Analysis and Zeta Functions*, Springer, 1977.

It contains the basic concepts including  $p$ -adic integration.

Of course you also find the basics in other books like

[Ba] Bachmann, G., *Introduction to  $p$ -adic Numbers and Valuation Theory*, Academic Press, New York 1964.

[Wa] Van der Waerden, B.L., *Algebra II*, Springer, 1967.

Adèles can be found in

[Ta] Tamagawa, T., Adèles, *Proc. Symp. Pure Math.* **9** (1966) 113–121, Amer. Math. Soc., Providence.

[Wei] Weil, A., *Basic Number Theory (3rd rev.ed.)*, Grundle Math. Wiss. 144, Springer, 1974.

Concerning the more speculative aspects of application of  $p$ -adic numbers to string theory, see

- [Man] Manin, Y.I., Reflections on Arithmetical Physics, Talk at Poiana-Brasov School on Strings and Conformal Field Theory, 1–14. September. 1987, (appeared in) *Perspect. Phys.*, Academic Press, Boston, MA, 1989.

Since the first edition of this book,  $p$ -adic and adelic mathematical physics developed further in the context of  $p$ -adic string theory and cosmology. For these more recent developments see

- [BF] Brekke, L., Freund, P.G.O.,  $p$ -adic Numbers in Physics, *Phys. Rep.* **233** (1993) 1–66.
- [DDNV] Djordjević, G.S., Dragovich, B., Nesić, L., Volovich, I.V.,  $p$ -adic and Adelic Minisuperspace Quantum Cosmology, *Int. J. Mod. Phys.*, A **17** (2002) 1413–1433.
- [KRV] Khrennikov, A.Y., Racić, Z., Volovich I.V. (eds),  $p$ -adic Mathematical Physics, *2nd International Conference*, Belgrade, Serbia, and Montenegro, 15–21.9.2005, AIP, Melville, New York, 2006.

---

# Index

- Abel's theorem, 58
- abelian varieties, 69
- absolute value, 203
- adèles, 209
- adelic string theory, 84
- adjunction formula, 189
- affine scheme, 160
- affine scheme associated to a  
    ring, 156
- affine space, 134
- affine variety, 95
- algebraic bundles, 91
- algebraic genus, 71
- algebraic map, 64
- algebraic set, 134
- algebraic space, 167
- algebraic stack, 167
- algebraic variety, 63
- analytic isomorphism, 14
- analyticmap, 14
- annulator, 151
- annulator ideal, 147
- arithmetic structure, 85
- associated sheaf, 119
- atlas, 7
  
- Batyrev's results, 193
- Belavin-Knizhnik theorem, 123
- Betti number, 24, 176
- biholomorphic map, 14
- Bogomolov-Tian-Todorov-Ran theorem,  
    194
- boundaries, 23
  
- Calabi-Yau manifold, 183
- canonical bundle, 107, 183
- canonical bundle of  $\mathbb{P}^n$ , 188
- canonical divisor, 36, 47
- canonical divisor class, 47, 100
- canonical homology bases, 28
- canonical line bundle, 100
- canonical sheaf, 100
- Cauchy-Riemann differential equations,  
    10
- Cech cocycles, 95
- Cech cohomology, 95
- Cech cohomology groups, 97
- chains, 22
- Chern character, 127
- Chern classes, 126
- Chern classes of a manifold, 183
- Chern number, 99
- Chow ring, 125
- Chow's theorem, 65
- closed differential forms, 49
- closed sets of  $\mathbb{P}^1$ , 62
- coarse moduli space, 73
- cocycle conditions, 88
- coherent sheaves, 94
- cohomologous cocycles, 89
- cohomology of a complex, 175
- cohomology sequence, 98
- compact manifold, 8
- compact Riemann surface, 13
- compactification, 79
- complete intersections, 191
- complete prime ideals, 150

- complex, 174
- complex manifold, 10
- complex projective space, 10
- complex torus, 185
- connected manifold, 7
- construction of the complex numbers, 204
- construction of the real numbers, 203
- coordinate change map, 8
- coordinate patch, 7
- cotangent bundle of the moduli space, 122
- covering, 28, 184
- covering group, 184
- covering transformations, 29
- curve, 64
- cuspidal cubic, 81
- cycles, 23
  
- deBar operator, 174
- deformations of complex structure, 194
- deformations of Kähler structure, 195
- degree of a divisor, 36
- degree of a hypersurface, 188
- degree of a line bundle, 108
- deRham operator, 174
- deRham cohomology group, 49, 175
- deRham differential, 174
- determinant bundle, 90
- diffeomorphism, 13
- differentiable isomorphism, 13
- differentiable manifold, 8
- differential, 49
- differential forms, 44
- differential forms of second order, 48
- dimension, 7
- dimension of a variety, 64
- direct image sheaf, 119
- discriminant function, 40
- divisor, 35, 64
- divisor class, 36
- Dolbeault cohomology group, 175
- Dolbeault differential, 174
- Dolbeault operator, 174
- double points, 80
- doubly periodic functions, 38
  
- Eisenstein series, 40
- elliptic curve, 66, 76, 185
- elliptic functions, 38
- elliptic modular function, 40
- Euler characteristic, 26, 176, 190
- Euler number, 176
- exact complex, 175
- exact differential forms, 49
- exponential sequence, 99
- exterior differential, 174
- exterior differentiation, 49
  
- family of Riemann surfaces, 71
- family of curves, 71
- field of meromorphic functions on the Riemann sphere, 34
- field of meromorphic functions, 32
- field of meromorphic functions on the torus, 40
- fine moduli space, 74
- finite unramified covering, 184
- fractional linear transformations, 77
- Fubini-Study form, 177
- functor of points, 161
- fundamental group, 18, 184
- fundamental region, 75
  
- Galois group, 149
- Gauss-Bonnet formula, 190
- genus, 13, 24
- geometric invariant theory, 80
- geometric mirror conjecture, 192
- geometric mirror symmetry, 170
- Givental's results, 200
- glueing function, 8
- Gromov-Witten invariants, 198
- Grothendieck group, 125
- Grothendieck-Riemann-Roch theorem, 128
  
- handle model, 26
- harmonic, 50
- Hausdorff space, 7
- higher dimensional tori, 53
- higher direct image sheaves, 119
- Hilbertscher Basissatz, 135
- Hilbertscher Nullstellensatz, 141
- Hirzebruch-Riemann-Roch theorem, 130

- Hodge bundle, 83, 122
- Hodge class, 82
- Hodge diamond, 180
- Hodge diamond of a Calabi-Yau 3-fold, 186
- Hodge diamond of K3 surfaces, 191
- Hodge diamond of the two-dimensional torus, 191
- Hodge number, 176
- Hodge numbers of the projective space, 181
- holomorphic bundles, 91
- holomorphic differential forms, 46
- holomorphic function, 32
- holomorphic vector fields on  $\mathbb{P}^1$ , 109
- holomorphic vector fields on higher genus Riemann surfaces, 111
- holomorphic vector fields on the torus, 111
- holonomy group, 187
- homeomorphism, 13
- homogeneous coordinates, 10
- homogeneous polynomials, 61
- homological mirror conjecture, 170, 195
- homology group, 23
- homotopy, 17
- hyperplane bundle, 188
- hypersurface, 136, 187
  
- idèles, 209
- ideal, 134
- integral domain, 141
- integration, 50
- intersection pairing, 57
- intersection product, 27
- irreducible, 63
- irreducible set, 137
- isothermal coordinates, 12
  
- $j$  function, 40
- Jacobi map, 57
- Jacobi's theorem, 58
- Jacobian, 56
  
- K3 surface, 186, 187
- Kähler manifold, 177
- Kähler cone, 179
- Kähler form, 177
- Kodaira embedding theorem, 65
  
- Krichever-Novikov (vector field) algebra, 115
- Krull dimension, 143
  
- large complex limit points, 195
- large Kähler limit points, 195
- Lefschetz hyperplane theorem, 189
- Lie algebra  $sl(2, \mathbb{C})$ , 111
- line bundle, 88
- linear affine subspace, 136
- linear variety, 63
- linearly equivalent, 36
- local-global principle, 209
- localization of a ring, 157
- locally free sheaves, 94
- locally ringed spaces, 160
- loop, 17
  
- manifold of genus 2, 25
- manifolds, 7
- Manin plane, 152
- mapping class group, 77
- maximal ideal, 139, 140, 150
- meromorphic differential forms, 46
- meromorphic function, 32
- meromorphic section of a line bundle, 103
- mirror pair, 192
- mirror symmetry, 169
- modular form, 84
- modular function, 75
- moduli space, 71
- moduli space of genus 1 curves, 74
- moduli space of higher genus curves, 77
- morphism of schemes, 160
- multiplicity, 33
- Mumford - Deligne - Knudsen compactification, 80
- Mumford isomorphism, 122, 129
  
- nil radical, 140
- nodal cubic, 81
- nodes, 80
- Noether formula, 190
- noetherian ring, 135
- non-archimedean valuation, 206
- nonsingular variety, 65
- number field, 211

- numbers of rational curves on a
  - quintic, 200
- numerical mirror conjecture, 196
- numerical mirror symmetry, 170
- orbifolds, 169
- order, 33, 35
- orientable manifold, 11
- Ortsuniformisierende, 31
- Ostrowski's theorem, 207
- $\mathbb{P}^n$ , 10
- $p$ -adic exponential series, 208
- $p$ -adic integers, 208
- $p$ -adic logarithm series, 208
- $p$ -adic numbers, 207
- $p$ -adic order, 208
- $p$ -adic string theory, 84, 211
- paracompact, 7, 8
- partition of unity, 8
- path, 17
- period matrix, 56
- Picard group, 36, 58, 82
- Picard group of the moduli functor, 83
- Picard variety, 58
- Poincaré duality, 176
- point derivation, 43
- polarization, 53
- Polyakov integration measure, 123
- polyhedron, 193
- presheaf, 119
- prime ideal, 140, 150
- primitive ideal, 151
- principal divisors, 35
- principal ideal rings, 135
- principally polarized torus, 53
- projective scheme, 164
- projective space, 164
- projective variety, 63
- proper algebraic map, 125
- $q$ -expansion, 76
- quadratic differentials, 79, 122
- quantum group, 152
- quantum plane, 152
- quasiprojective variety, 63
- quintics in  $\mathbb{P}^4$ , 196
- radical ideal, 140
- radical of an ideal, 140
- ramified covering, 28
- rational functions, 64
- real analytic manifold, 8
- reduced, 140
- representable functor, 161
- residue field, 158
- residue theorem, 46
- restricted holonomy group, 187
- Ricci flat, 184
- Riemann sphere, 11, 48
- Riemann surface, 13
- Riemann zeta function, 210
- Riemann–Roch theorem, 36
- Riemann–Roch theorem, 129
- Riemann–Roch theorem for line
  - bundles, 108
- ring of fractions, 156
- $S^n$ , 8
- Satake compactification, 79
- scheme, 159
- Schottky problem, 78
- sections of a vector bundle, 89
- self intersection number, 190
- Serre duality, 79, 100, 121, 176
- sheaf cohomology theory, 95
- sheaf homomorphism, 92
- sheaf of  $\mathcal{O}$ -modules, 92
- sheaf of relative differentials, 120
- sheaf, sheaves, 91
- short exact sequence of sheaves, 93
- Siegel's upper half-space, 67, 78
- simple module, 147
- simplices, 21
- simplicial homology group, 23
- simply connected, 19
- singular homology, 30
- singular point, 65
- singular variety, 65
- smooth variety, 65
- sphere, 8, 25
- stable curves, 80
- standard coordinates, 31
- standard covering of the sphere, 9
- Stokes' Theorem, 51
- string compactification, 169
- structure sheaf, 92, 158

- surface, 64
- symplectic basis, 54
- symplectic group, 55, 59
  
- tangent bundle, 87
- tangent bundle of the moduli space, 122
- tangent space, 43
- Teichmüller space, 77
- Teichmüller surface, 77
- tensors, 90
- theta function, 67
- theta functions with characteristics, 68
- theta null values, 84
- Todd class, 127
- topological isomorphism, 13
- Torelli's theorem, 58, 78
- toric variety, 193
- torus, 25, 38, 48, 65, 178
- triangulation, 21
- trivializing covering, 88
- type of a differential form, 173
  
- ultrametric, 206
- universal covering, 28, 184
- universal family, 74
- upper half plane, 178
  
- valuation for a field, 205
- vanishing set of an ideal, 134
- variety, 62, 137
- vector bundle, 88
- very ample line bundle, 100
- Virasoro algebra, 117
  
- Weierstraß points, 113
- Weierstraß  $\wp$ -function, 39
- weighted projective space, 193
- Weil divisor, 64
- Wess-Zumino-Novikov-Witten models, 118
  
- Yukawa coupling, 200
  
- Zariski topology, 62, 136
- Zariski topology of a ring, 144

# Theoretical and Mathematical Physics

---

**An Introduction to Riemann Surfaces, Algebraic Curves and Moduli Spaces**

2nd enlarged edition

By M. Schlichenmaier

**Quantum Probability and Spectral Analysis of Graphs**

By A. Hora and N. Obata

**From Nucleons to Nucleus**

Concepts of Microscopic Nuclear Theory

By J. Suhonen

**Concepts and Results in Chaotic Dynamics: A Short Course**

By P. Collet and J.-P. Eckmann

**The Theory of Quark and Gluon Interactions**

4th Edition

By F. J. Ynduráin

**From Microphysics to Macrophysics**

Methods and Applications of Statistical Physics

Volume I, Study Edition

By R. Balian

---

**Titles published before 2006 in *Texts and Monographs in Physics***

**The Statistical Mechanics of Financial Markets**

3rd Edition

By J. Voit

**Magnetic Monopoles**

By Y. Shnir

**Coherent Dynamics of Complex Quantum Systems**

By V. M. Akulin

**Geometric Optics on Phase Space**

By K. B. Wolf

**General Relativity**

By N. Straumann

**Quantum Entropy and Its Use**

By M. Ohya and D. Petz

**Statistical Methods in Quantum Optics I**

By H. J. Carmichael

**Operator Algebras and Quantum Statistical Mechanics I**

By O. Bratteli and D. W. Robinson

**Operator Algebras and Quantum Statistical Mechanics 2**

By O. Bratteli and D. W. Robinson

**Aspects of Ergodic, Qualitative and Statistical Theory of Motion**

By G. Gallavotti, F. Bonetto and G. Gentile

**The Frenkel-Kontorova Model**

Concepts, Methods, and Applications

By O. M. Braun and Y. S. Kivshar

**The Atomic Nucleus as a Relativistic System**

By L. N. Savushkin and H. Toki

**The Geometric Phase in Quantum Systems**

Foundations, Mathematical Concepts, and Applications in Molecular and Condensed Matter Physics

By A. Bohm, A. Mostafazadeh, H. Koizumi,

Q. Niu and J. Zwanziger

**Relativistic Quantum Mechanics**

2nd Edition

By H. M. Pilkuhn

**Physics of Neutrinos**

and Applications to Astrophysics

By M. Fukugita and T. Yanagida

**High-Energy Particle Diffraction**

By E. Barone and V. Predazzi

**Foundations of Fluid Dynamics**

By G. Gallavotti

**Many-Body Problems and Quantum Field Theory An Introduction**

2nd Edition

By Ph. A. Martin, F. Rothen,

S. Goldfarb and S. Leach

**Statistical Physics of Fluids**

Basic Concepts and Applications

By V. I. Kalikmanov

**Statistical Mechanics A Short Treatise**

By G. Gallavotti



**Quantum Non-linear Sigma Models**

From Quantum Field Theory  
to Supersymmetry, Conformal Field Theory,  
Black Holes and Strings  
By S. V. Ketov

**Perturbative Quantum Electrodynamics and  
Axiomatic Field Theory**

By O. Steinmann

**The Nuclear Many-Body Problem**

By P. Ring and P. Schuck

**Magnetism and Superconductivity**

By L.-P. Lévy

**Information Theory and Quantum Physics**

Physical Foundations for Understanding the  
Conscious Process  
By H. S. Green

**Quantum Field Theory in Strongly  
Correlated Electronic Systems**

By N. Nagaosa

**Quantum Field Theory in Condensed Matter  
Physics**

By N. Nagaosa

**Conformal Invariance and Critical  
Phenomena**

By M. Henkel

**Statistical Mechanics of Lattice Systems**

Volume 1: Closed-Form and Exact Solutions  
2nd Edition  
By D. A. Lavis and G. M. Bell

**Statistical Mechanics of Lattice Systems**

Volume 2: Exact, Series  
and Renormalization Group Methods  
By D. A. Lavis and G. M. Bell

**Fields, Symmetries, and Quarks**

2nd Edition  
By U. Mosel

**Renormalization** An Introduction

By M. Salmhofer

**Multi-Hamiltonian Theory of Dynamical  
Systems**

By M. Błaszak

**Quantum Groups and Their Representations**

By A. Klimyk and K. Schmüdgen

**Quantum** The Quantum Theory of Particles,  
Fields, and Cosmology

By E. Elbaz

**Effective Lagrangians for the Standard Model**

By A. Dobado, A. Gómez-Nicola,  
A. L. Maroto and J. R. Peláez

**Scattering Theory of Classical  
and Quantum  $N$ -Particle Systems**

By. J. Dereziński and C. Gérard

**Quantum Relativity** A Synthesis

of the Ideas of Einstein and Heisenberg  
By D. R. Finkelstein

**The Mechanics and Thermodynamics  
of Continuous Media**

By M. Šilhavý

**Local Quantum Physics** Fields, Particles,

Algebras  
2nd Edition  
By R. Haag

**Relativistic Quantum Mechanics  
and Introduction to Field Theory**

By F. J. Ynduráin

**Supersymmetric Methods in Quantum  
and Statistical Physics**

By G. Junker

**Path Integral Approach  
to Quantum Physics** An Introduction

2nd printing  
By G. Roepstorff

**Finite Quantum Electrodynamics**

The Causal Approach  
2nd edition  
By G. Scharf

**From Electrostatics to Optics**

A Concise Electrodynamics Course  
By G. Scharf

**Geometry of the Standard Model  
of Elementary Particles**

By A. Derdziński

**Quantum Mechanics II**

By A. Galindo and P. Pascual

**Generalized Coherent States**

and Their Applications  
By A. Perelomov

**The Elements of Mechanics**

By G. Gallavotti

**Essential Relativity** Special, General,  
and Cosmological Revised

2nd edition  
By W. Rindler